

---

## Perspective In Ophthalmology

---

# There may be virus in conjunctival secretion of patients with COVID-19

Liang Liang  and Ping Wu

Department of Ophthalmology, Yichang Central People's Hospital, The first college of Clinical Medical Science, China Three Gorges University, Yichang, P. R. China

**Key words:** coronavirus – pneumonia – conjunctival – secretion

Acta Ophthalmol.

© 2020 Acta Ophthalmologica Scandinavica Foundation. Published by John Wiley & Sons Ltd

doi: 10.1111/aos.14413

The COVID-19 is a public health emergency. The COVID-19 patient is the main source of infection, and asymptomatic infective patient can also be a source of infection (Bhadelia 2020). The main route of transmission is through respiratory droplets and contact (Burki 2020).

It is not clear whether the 2019-nCoV is transmitted through the mucous membrane of the eye. S-protein of SARS-like coronavirus can interact with human ACE2 protein and infect human respiratory epithelial cells (Ge et al. 2013). The human cornea and conjunctiva express ACE2 receptor, which can theoretically bind to the 2019-nCoV and cause infection. Our hospital is from the epidemic area. In our hospital, 37 patients of 2019-

nCoV infection pneumonia were detected the nucleic acid in the conjunctival sac by real-time RT-PCR. According the Chinese COVID-19 diagnosis and treatment (V) (Huang et al. 2020), 12 cases were severe patients, and the others were mild patients. Three cases had conjunctival congestion and other inflammatory appearance. One case of severe patients, which conjunctival sac secretion nucleic acid test was positive by real-time RT-PCR, but this severe patient had no conjunctivitis. The other 36 patients were negative in nucleic acid test of conjunctival secretion.

Therefore, we must pay attention to the possibility of virus in conjunctival sac secretion of COVID-19 patients and need to further detect the virus in conjunctival sac secretion as evidence. The viral load of conjunctival sac secretion of COVID-19 patients is relatively low, and we estimate the viral load is directly proportional to the severity of the disease. Whether the 2019-nCoV can be transmitted through conjunctiva is further studied.

## References

- Bhadelia N (2020): Coronavirus: hospitals must learn from past pandemics. *Nature* **578**: 193–193.
- Burki TK (2020): Coronavirus in China. *Lancet Respir Med* **8**: 238.
- Ge XY, Li JL, Yang XL et al. (2013): Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* **503**: 535–8.
- Huang C, Wang Y, Li X et al. (2020): Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**: 497–506.

---

Received on February 27th, 2020.

Accepted on March 1st, 2020.

### Correspondence:

Liang Liang  
Department of Ophthalmology  
Yichang Central People's Hospital  
The first college of Clinical Medical Science  
China Three Gorges University  
Yichang 443003  
Hubei  
P. R. China  
Tel: 08607176481665  
Fax: 08607176482302  
Email: liangliang419519@163.com